

HER number

131

Identification

<i>Name</i> Tb	<i>Morphotype</i> C1 (Podophage)	<i>Other designations</i> TB, Tbilisi	
-------------------	-------------------------------------	--	--

Taxonomy

<i>Realm</i> <i>Duplodnaviria</i>	<i>Kingdom</i> <i>Heunggongvirae</i>	<i>Phylum</i> Uroviricota	<i>Class</i> <i>Caudoviricetes</i>
<i>Order</i>	<i>Family</i>	<i>Genus</i> <i>Perisivirus</i>	<i>Species</i> <i>Perisivirus Tb</i>

Images

Electron Micrograph

Image

FHRCEV




Image description

Magnification: 297,000X

Bar: 50 nm

Staining: PTA

Characteristics

Plaques: 0.5- 1mm, clear, sloping borders.
Host is class III pathogen.
Albimi broth, Serum dextrose agar.
Growth time = 36 h.

Genomic sequence

Activated

Propagation conditions

Bacterial hosts

1131

Reference

Brinley-Morgan, W. J. D. Kay, and D. E. Bradley. 1960. *Brucella* bacteriophage. Nature (London). 188:74-75.

Remarks

History

History

Received from

M. J. Corbel,
FAO/WHO Collaborating Centre for Reference and
Research on Brucellosis,
Ministry of Agriculture, Fisheries and Food,
Central Veterinary Laboratory,
New Haw, Weybridge, Surrey, England,
KT15 3NB

Date

06-22-1982

Isolated by

M. Z. Popkhadze, and T. G. Abashidze.
In Tiflis, Georgia,
URSS

Date

Source

Liquid manure of cowshed.

Updated at
2024-01-16