

HER number

192

Identification

Name 192	Morphotype B1 (Siphophage)	Other designations	

Taxonomy

Realm	Kingdom	Phylum	Class
Duplodnaviria	Heunggongvirae	<u>Uroviricota</u>	Caudoviricetes
Order	Family	Genus	Species

Images

Electron Micrograph

H.-W.ACKERMANN

Image description

Magnification: 297,000X

Bar: 50 nm Staining: UA

Characteristics

Plaques: 1 mm, clear with halo.

Genome of 63.8 kb.

Occurs in temperate (LP52) and lytic (LP52c)

forms.

Restriction map is identical with that of phage

LP51.

Genomic DNA shows 50% cross-hybridization with the DNA of phage c.

Genomic sequence

Deactivated

Propagation conditions

Bacterial hosts

1190

Reference

Thorne, C.B., and B.B. Kowalski. 1976. Temperate bacteriophages for *Bacillus licheniformis*. In Microbiology, edited by D. Schlessinger, American Society for Microbiology, Washington D.C., p. 302-314.

Imanaka, T., K. Uchida, M. Tateishi, S. Aiba. 1979. Inducible bacteriophages of *Bacillus licheniformis* ATCC 10716. Virology 95:249-250.

Remarks

History

History

Received from

Dr. J. Doskocil Institute of Molecular Genetics

Flemingovo nàm. 2, 16637 Praha 6,

Czechoslovakia

Received from

Prof. Curtis B. Thorne Department of Microbiology University of Massachusetts Amhersy, Massachusetts 01002, USA

Date

Date

20-09-1983

Isolated by

Prof. Curtis B. Thorne

University of Massachusetts, Amherst,

USA

Date 1976

Source

Bacterial strain 10716 induced with mitomycin C (0.5mg/ml)

Updated at

2024-01-16