

HER  
1203

## Identification

|                                    |                                 |                                      |               |                |
|------------------------------------|---------------------------------|--------------------------------------|---------------|----------------|
| <i>Genus</i><br><i>Lactococcus</i> | <i>Species</i><br><i>lactis</i> | <i>Subspecies</i><br><i>cremoris</i> | <i>Biovar</i> | <i>Serovar</i> |
| <i>Strain</i><br>158               |                                 | <i>Aliases</i>                       |               |                |

## Growth conditions

|                    |                       |                          |                         |  |
|--------------------|-----------------------|--------------------------|-------------------------|--|
| <i>Temp.</i><br>30 | <i>Medium</i><br>GM17 | <i>Aerobic/Anaerobic</i> | <i>Agitation</i><br>Non | <i>Other</i><br>10mM CaCl <sub>2</sub> |
|--------------------|-----------------------|--------------------------|-------------------------|--|

## Lysogeny

## Phage sensitivity

HER 203 (936)

## Reference

### Remark

Cells in liquid medium must be washed with fresh medium before the inoculation of a new liquid medium. The pH of M17 medium is very important.

## History

### History

#### Received from

Dr A.W. Jarvis, New Zealand Dairy Research Institute,  
Private Bag, Palmerston North, New Zealand

#### Date

04-02-1984

*Last update*

2016-09-28