Host HER 1237

HER
1237

Identification

Genus	*Species*	*Subspecies*	*Biovar*	*Serovar*
Halobacterium	*salinarum*			

Strain	*Aliases*
NRC 34001	

Growth conditions

Temp.	*Medium*	*Aerobic/Anaerobic*	*Agitation*	*Other*
37	3.5S			

Lysogeny

Phage sensitivity

HER 237 (S45)

HER 343 (Ja.1)

Reference
Lockhead, A.G. 1934. Can. J. Res. 10: 275-286.

Remark
Also held in the NRC permament collection (no. 34001) and in the ATCC collection (no. 33170)
For reproducible results, it is best to maintain the organism in log phase in broth, checking frequently that the cells demonstrate regular morphology and appear as slender rods.
Log phase cultures can be refrigerated for 2-3 weeks.

History

History

Received from	**Date**
Dr Allen C. Wais, Department of Biochemistry, Faculty of Medecine, U.W.I. Mona, Kingston 7, Jamaica, West Indies	3-22-1985
Received from	**Date**
R.E. MacDonald, Cornell University, Ithaca, NY, USA	02-1973
Received from	**Date**
D.J. Kushner, Dept. Of Biology, University of Ottawa	
Isolated by	**Date**
A.G. Lockhead, Division of Bacteriology, Dominion Experimental Farms, Ottawa, Canada.	1934

Last update

2012-02-24