

HER  
1237

## Identification

<i>Genus</i> <i>Halobacterium</i>	<i>Species</i> <i>salinarum</i>	<i>Subspecies</i>	<i>Biovar</i>	<i>Serovar</i>
<i>Strain</i> NRC 34001		<i>Aliases</i>		

## Growth conditions

<i>Temp.</i> 37	<i>Medium</i> 3.5S	<i>Aerobic/Anaerobic</i>	<i>Agitation</i>	<i>Other</i>
--------------------	-----------------------	--------------------------	------------------	--------------

## Lysogeny

## Phage sensitivity

HER 237 (S45)

HER 343 (Ja.1)

### Reference

Lockhead, A.G. 1934. Can. J. Res. 10: 275-286.

### Remark

Also held in the NRC permanent collection (no. 34001) and in the ATCC collection (no. 33170)

For reproducible results, it is best to maintain the organism in log phase in broth, checking frequently that the cells demonstrate regular morphology and appear as slender rods. Log phase cultures can be refrigerated for 2-3 weeks.

## History

*History*

**Received from**

Dr Allen C. Wais, Department of Biochemistry, Faculty of Medicine, U.W.I. Mona, Kingston 7, Jamaica, West Indies

**Date**

3-22-1985

**Received from**

R.E. MacDonald, Cornell University, Ithaca, NY, USA

**Date**

02-1973

**Received from**

D.J. Kushner, Dept. Of Biology, University of Ottawa

**Date**

**Isolated by**

A.G. Lockhead, Division of Bacteriology, Dominion Experimental Farms, Ottawa, Canada.

**Date**

1934

*Last update*

2012-02-24