

HER
1307**Identification**

<i>Genus</i> Vibrio	<i>Species</i> parahaemolyticus	<i>Subspecies</i>	<i>Strain</i> 16
------------------------	------------------------------------	-------------------	---------------------

*Aliases***Growth conditions**

<i>Temp.</i> 25	<i>Medium</i> NaCl	<i>Aerobic/Anaerobic</i>	<i>Agitation</i>	<i>Other</i>
--------------------	-----------------------	--------------------------	------------------	--------------

Lysogeny**Phage sensitivity**

HER 307 (?16)	HER 308 (?HAWI-5)	HER 309 (?PEL8C-1)
---------------	-------------------	--------------------

Reference

Kellogg, C.A., J.B. Rose, S.C. Jiang, J.M. Thurmond, and J.H. Paul. 1995. Genetic diversity of related vibriophages isolated from marine environments around Florida and Hawaii, USA. Marine Ecol. Progr. Ser. 120:89-98.

Remark

Some natural resistance to nalidixic acid.
Has photoreactivation and nucleotide excision repair of UV damage.
Used to isolate over 200 phages from marine environments around Florida, the Gulf of Mexico, Hawaii.

History*History***Received from**

Dr Christina A. Kellogg, University of South Florida, Marine Science Department, 140 7th Avenue S., St. Petersburg, FL33701, USA

Date

09-22-1998

Last update

2012-02-29