

HER  
1307

## Identification

<i>Genus</i>	<i>Species</i>	<i>Subspecies</i>	<i>Biovar</i>	<i>Serovar</i>
<i>Vibrio</i>	<i>parahaemolyticus</i>			
<i>Strain</i>		<i>Aliases</i>		
16				

## Growth conditions

<i>Temp.</i>	<i>Medium</i>	<i>Aerobic/Anaerobic</i>	<i>Agitation</i>	<i>Other</i>
25	NaCl			

## Lysogeny

## Phage sensitivity

HER 307 (φ16)

HER 308 (φHAWI-5)

HER 309 (φPEL8C-1)

### Reference

Kellogg, C.A., J.B. Rose, S.C. Jiang, J.M. Thurmond, and J.H. Paul. 1995. Genetic diversity of related vibriophages isolated from marine environments around Florida and Hawaii, USA. *Marine Ecol. Progr. Ser.* 120:89-98.

### Remark

Some natural resistance to nalidixic acid.

Has photoreactivation and nucleotide excision repair of UV damage.

Used to isolate over 200 phages from marine environments around Florida, the Gulf of Mexico, Hawaii.

## History

*History*

**Received from**

Dr Christina A. Kellogg, University of South Florida,  
Marine Science Department, 140 7th Avenue S., St.  
Petersburg, FL33701, USA

**Date**

09-22-1998

*Last update*

2012-02-29