

*HER number*  
166

## Identification

<i>Name</i> 166	<i>Morphotype</i> B1 (Siphophage)	<i>Other designations</i>
--------------------	--------------------------------------	---------------------------


## Taxonomy

<i>Realm</i>	<i>Kingdom</i>	<i>Phylum</i>	<i>Class</i>
<i>Order</i>	<i>Family</i> ?1	<i>Genus</i>	<i>Species</i>

## Images

*Electron Micrograph*

*Image*



*Image description*

Magnification: 297,000X

Bar: 50 nm

Staining: PTB

<i>Characteristics</i> Plaques : 0.5 mm, veiled. Capsid covered with spherical projections. Relative of VP3.	<i>Genomic sequence</i> Deactivated
<b>Propagation conditions</b>	
<i>Bacterial hosts</i> 1165	
<i>Reference</i> Koga, T., S. Toyoshima, and T. Kawata. 1982. Morphological varieties and host ranges of *Vibrio parahaemolyticus* bacteriophages isolated from seawater. Appl. Environ. Microbiol. 44:466-470.	
<i>Remarks</i>	
<b>History</b>	
<i>History</i>	
<b>Isolated by</b> T. Koga and T. Kawata Tokushima, Japan	<b>Date</b> 06-1980
<b>Received from</b> Dr Tomio Kawata, Department of Food Microbiology, Tokushima University School of Medecine, Tokushima, 770, Japan	<b>Date</b> 21-02-1983
<i>Source</i> Seawater	
<i>Updated at</i> 2024-01-16	