

HER number
168

Identification

<i>Name</i> 168	<i>Morphotype</i> B1 (Siphophage)	<i>Other designations</i>
--------------------	--------------------------------------	---------------------------

Taxonomy

<i>Realm</i> <i>Duplodnaviria</i>	<i>Kingdom</i> <i>Heunggongvirae</i>	<i>Phylum</i> <i>Uroviricota</i>	<i>Class</i> <i>Caudoviricetes</i>
<i>Order</i>	<i>Family</i>	<i>Genus</i>	<i>Species</i>

Images

Electron Micrograph

Image




Image description

Magnification: 297,000X

Bar: 50 nm

Staining: UAB

<i>Characteristics</i> Plaques smaller than 0.1 mm, clear. Ressemble cholera phage IV, but has no tail fibers.	<i>Genomic sequence</i> Deactivated
Propagation conditions	
<i>Bacterial hosts</i> 1165	
<i>Reference</i> Koga, T., S. Toyoshima, and T. Kawata. 1982. Morphological varieties and host ranges of *Vibrio parahaemolyticus* bacteriophages isolated from seawater. Appl. Environ. Microbiol. 44:466-470.	
<i>Remarks</i>	
History	
<i>History</i>	
Received from Dr Tomio Kawata, Department of Food Microbiology, Tokushima University School of Medecine, Tokushima, 770, Japan	Date 02-21-1983
Isolated by T. Koga and T. Kawata Tokushima, Japan	Date 07-1980
<i>Source</i> Seawater	
<i>Updated at</i> 2024-01-16	