

*HER number*  
169

## Identification

<i>Name</i> 169	<i>Morphotype</i> B1 (Siphophage)	<i>Other designations</i>	
--------------------	--------------------------------------	---------------------------	--

## Taxonomy

<i>Realm</i> <i>Duplodnaviria</i>	<i>Kingdom</i> <i>Heunggongvirae</i>	<i>Phylum</i> <a href="#"><i>Uroviricota</i></a>	<i>Class</i> <i>Caudoviricetes</i>
<i>Order</i>	<i>Family</i>	<i>Genus</i> <i>Enhodamvirus</i>	<i>Species</i> <i>Enhodamvirus VP5</i>

## Images

*Electron Micrograph*

*Image*  
FHRCEV



*Image description*

Magnification: 297,000X

Bar: 50 nm

Staining: UA

<i>Characteristics</i>  Plaques of 0.5 mm, extremely opaque, more visible after 3 days. Elongated head.	<i>Genomic sequence</i> Activated
<b>Propagation conditions</b>	
<i>Bacterial hosts</i> 1169	
<i>Reference</i> Koga, T., S. Toyoshima, and T. Kawata. 1982. Morphological varieties and host ranges of *Vibrio parahaemolyticus* bacteriophages isolated from seawater. Appl. Environ. Microbiol. 44:466-470.	
<i>Remarks</i>	
<b>History</b>	
<i>History</i>	
<b>Received from</b> Dr Tomio Kawata, Department of Food Microbiology, Tokushima University School of Medecine, Tokushima, 770, Japan	<b>Date</b> 02-21-1983
<b>Isolated by</b> T. Toga and T. Kawata Tokushima, Japan	<b>Date</b> 07-1979
<i>Source</i> Seawater	
<i>Updated at</i> 2024-01-16	