

HER number
175

Identification

| | | |
|--------------------|--------------------------------------|-------------------------------------|
| <i>Name</i> 175 | <i>Morphotype</i> B1 (Siphophage) | <i>Other designations</i> SPbeta |
|--------------------|--------------------------------------|-------------------------------------|

Taxonomy

| | | | |
|--------------|---------------------|-----------------------------|--------------------------------------|
| <i>Realm</i> | <i>Kingdom</i> | <i>Phylum</i> | <i>Class</i> |
| <i>Order</i> | <i>Family</i> ?1 | <i>Genus</i> Spbetavirus | <i>Species</i> Spbetavirus SPbeta |

Images

Electron Micrograph

Image



Image description

Magnification: 297,000X

Bar: 50 nm

Staining: UAB

Characteristics
Plaques: 0.2-0.5 mm, clear.

Genomic sequence
Activated

Propagation conditions

Bacterial hosts

1175

Reference

Warner, F.D., G.A. Kitos, M.P. Romano, and H.E. Hemphill. 1977. Characterization of SP12 : a temperate bacteriophage from *Bacillus subtilis* 168M. Can. J. Microbiol. 23:45-51.

Remarks

Held in another permanent collection : Bacillus Genetic Stock Center
May be obtained from strain HER 1171 by induction with mitomycin C (1mg/ml)
First observed in *B. subtilis* lysates before 1963.

History

History

Received from

H.E. Hemphill,
Department of Microbiology,
School of Medicine,
University of Washington,
Seattle, Washington 98195,
USA

Date

Received from

S.A. Zahler,
Section of Genetics and Development,
Bradfield Hall,
Cornell University,
Ithaca, NY 14853,
USA

Date

31-03-1983

Isolated by

F.D. Warner et al.,
Department of Biology,
Syracuse University,
Syracuse, NY 13210,
USA

Date

Before 1977

Source

Bacillus subtilis 168M

Updated at

2024-01-16