

Numéro HER
191

Identification

| | | | |
|-----------|-------------------------------|----------------------------|--|
| Nom θc | Morphotype B3 (Siphophage) | Autres désignations BLE | |
|-----------|-------------------------------|----------------------------|--|

Taxonomie

| | | | |
|---------------------------------|--------------------------------|--|---------------------------------|
| Domaine <i>Duplodnaviria</i> | Règne <i>Heunggongvirae</i> | Embranchement Uroviricota | Classe <i>Caudoviricetes</i> |
| Ordre | Famille | Genre | Espèce |

Images

| | |
|--|---------------------------------|
| Photo en microscopie électronique | Description de l'image |
| Caractéristique Plaques: 1.0 mm, clear. | Séquence génomique Désactivé |

Conditions de propagation

Hôtes bactériens
1190

Référence

Doskocil, J., J. Forstova, and J. Stokrova. 1978. Temperate and virulent forms of phage θc attacking *Bacillus licheniformis*. Mol. Gen. Genet. 160:311-317.

Remarque

Phage BLE was isolated from the same batch independantly by Dr Ludvik:
Ludvik, J., L. Erbenova, and H. Lipavska. 1977. Ultrastructure of *Bacillus licheniformis*
bacteriophage BLE and its DNA. *Virology* 77:872-875.

Historique

Historique

Received from

Dr. J. Dosekocil
Institute of Molecular Genetics
Flemingovo nám. 2,
16637 Praha 6,
Czechoslovakia

Date

09-20-1983

Isolated by

J. Dosekocil
Kourim,
Czechoslovakia

Date

1975

Source

A lysed industrial batch of *B. licheniformis* 10716.

Dernière mise à jour

2024-01-16