

Numéro HER
317

Identification

Nom 33D (Warsaw)	Morphotype B1 (Siphophage)	Autres désignations
---------------------	-------------------------------	---------------------

Taxonomie

Domaine <i>Duplodnaviria</i>	Règne <i>Heunggongvirae</i>	Embranchement Uroviricota	Classe <i>Caudoviricetes</i>
Ordre	Famille	Genre	Espèce

Images

Photo en microscopie électronique	Description de l'image
Caractéristique Clear plaques of 1.0 mm	Séquence génomique Désactivé

Conditions de propagation

Hôtes bactériens
1316

Référence
Jones, W.D. 1975. Differentiation of known strains of BCG from isolates of *Mycobacterium bovis* and *Mycobacterium tuberculosis* by using mycobacteriophage 33D. J. Clin. Microbiol. 1:391-392

Remarque

This phage lyses *M. tuberculosis* and wild strains of *M. bovis*; it does not lyse any of the attenuated strains of BCG.

Used in clinical settings for differentiation of BCG from *M. bovis* and *M. tuberculosis*. (1-RTD on ATCC 607)

Historique

Historique

Received from

Dr Edith Mankiewicz
Lakeshore General Hospital
160 Stillview
Pointe-Claire, QC, Canada
H9R 2Y2

Date

04-14-1988

Received from

Dr M. Buraczewska

Date**Isolated by**

Dr M. Buraczewska
Watsaw, Poland

Date

1970

Source

Lysogenic bacilli

Dernière mise à jour

2024-01-19